

AD-777 901

MAMMALIAN TOXICOLOGY AND TOXICITY TO
AQUATIC ORGANISMS OF WHITE PHOSPHORUS
AND "PHOSSY WATER" A WATERBORNE
MUNITIONS MANUFACTURING WASTE POLLU-
TANT - A LITERATURE EVALUATION

Dickinson Burrows, et al

Associated Water and Air Resources Engineers,
Incorporated

Prepared for:

Army Medical Research and Development
Command

November 1973

DISTRIBUTED BY:

NTIS

National Technical Information Service
U. S. DEPARTMENT OF COMMERCE
5285 Port Royal Road, Springfield Va. 22151

ACKNOWLEDGMENTS

The authors wish to acknowledge the conscientious efforts of Mr. Robert C. Kahn and Miss Sandra L. Gordon in the acquisition and assembly of information used in this report. They also wish to acknowledge the technical guidance and editorial efforts of Dr. David H. Rosenblatt of the Environmental Quality Division, US Army Medical Bioengineering Research and Development Laboratory.

TABLE OF CONTENTS

ABSTRACT	i
ACKNOWLEDGMENTS	11
TABLE OF CONTENTS	111
LIST OF TABLES AND FIGURES	v
LIST OF ABBREVIATIONS	vi
INTRODUCTION	1
Background	1
Approach to Mammalian Toxicology	2
Approach to Toxicity to Aquatic Organisms	2
Current Research on WP Toxicology and "Phossy Water"	3
MAMMALIAN TOXICOLOGY OF WHITE PHOSPHORUS AND RELATED MATERIALS	3
Acute Toxicity in Animals	3
Acute Toxicity in Man	3
Chronic Toxicity in Animals	5
Chronic Toxicity in Man (Including Potential for Carcinogenicity, Mutagenicity, and Teratogenicity)	5
Burns due to Phosphorus	6
Toxicity of Phosphine	6
BIOCHEMICAL STUDIES	8
TOXICITIES OF WHITE PHOSPHORUS TO AQUATIC ORGANISMS	10
Toxicity of Elemental Phosphorus in Water	10
Accumulation of Phosphorus by Aquatic Organisms	11
Toxicity to Waterfowl	13
Toxicity of the Total Waste	13
Behavioral Effects	19
Pathology	19
Summary and Conclusions	20
CHEMISTRY OF WHITE PHOSPHORUS AND DERIVED SUBSTANCES	21
The Fate of Elemental Phosphorus in Water	21

ANALYTICAL METHODOLOGY	21
Elemental Phosphorus	21
Phosphine	22
Oxyacids of Phosphorus	22
RECOMMENDATIONS FOR FUTURE EXPERIMENTAL AND FIELD STUDIES ON WHITE PHOSPHORUS	23
LITERATURE CITED	25
APPENDIX A Sources Consulted	38
APPENDIX B Uncited References	41

LIST OF TABLES AND FIGURES

- TABLE 1 Summary of Acute Toxicity of White Phosphorus in Animals
- TABLE 2 Summary of Chronic Toxicity of White Phosphorus to Humans (Heimann, 1946)⁴
- TABLE 3 Summary of the Physical Response to Various Concentrations of Phosphine (Henderson and Haggard, 1943)⁷¹
- TABLE 4 Summary of Toxicity of White Phosphorus to Fish Species and Crustaceans
- TABLE 5 Phosphorus Accumulation by Seaweed and Invertebrates (Fletcher, 1971)⁹⁰
- TABLE 6 Enzyme Inhibition by Phosphite and Orthophosphite
- TABLE 7 Analysis of Long Harbour Wastewater (Idler, 1969)⁵
- FIGURE 1 Toxicities of Various Substances to Seawater Maintained Brook Trout (Fletcher *et al.*, 1971)⁸⁹

LIST OF ABBREVIATIONS

AAP	Army Ammunition Plant
ACGIH	American Conference of Government Industrial Hygienists
ca	circa (about)
CN	Cyanide
DOD	Department of Defense
g	gram
gpm	gallons per minute
kg	kilogram
iv	intravenous
LC50	Lethal Concentration (kills 50%)
LD	Lethal Dose
LD50	Lethal Dose (kills 50%)
LT	Lethal Time
LT50	Lethal Time (kills 50%)
M	Molar
MAC	Maximum Allowable Concentration
mg	milligram
MGD	Million Gallons per Day
or	oral
P, P ₄	Phosphorus
³² P	Phosphorus isotope 32
ppb	parts per billion
ppm	parts per million
sc	subcutaneous
TLV	Threshold Limit Value
WP	White Phosphorus
ug	micro-gram

INTRODUCTION

The objective of this work is to provide part of a data base for standards for military-unique product discharges into waters subject to State and Federal regulations.

Background. This report is concerned with the toxicology of elemental "white" phosphorus (also known as "yellow" phosphorus) and its probable aqueous decomposition products. The generation of "phossy water" wastes in loading operations at the Pine Bluff, Arkansas, Army Ammunition Plant and some environmental consequences thereof have been reviewed by Rosenblatt, Small and Barkley (1973)¹. White phosphorus (WP) is purchased commercially and unloaded at Pine Bluff Arsenal. It is shipped in rail tank cars and transferred molten, under a warm water blanket, to munitions loading operations.

"Phossy water" is defined as water that has come in direct contact with large quantities of molten phosphorus, including dip-tank overflow, water used to flush white phosphorus from pipes, spray water used for wetting the dip-fill line conveyor and water from periodic cleaning of the dip-fill tanks. The rate of production of this phosphorus-laden water varies greatly, as does the white phosphorus concentration in the water. Elemental phosphorus is continually undergoing reaction with dissolved oxygen (under water), or with atmospheric oxygen to produce oxides of phosphorus that hydrolyze to give phosphorus compounds, perhaps in various oxidation states. Given adequate time, the white phosphorus particulates settle out as a sludge and the dissolved phosphorus is slowly oxidized. However, streams containing white phosphorus remain noticeably polluted over considerable distances.

Acute phosphorus poisoning early in this century was caused by the ingestion of match tips, fireworks, and quack nostrums, and is still encountered with surprising regularity. The literature abounds in cases of human poisonings after the ingestion of rat poisons or roach powders where phosphorus may be present in concentrations up to 4%. The conquest of occupational phosphorus poisoning in lucifer match manufacturing has been described by Oliver (1938)² and by Hamilton and Hardy (1949)³.

Chronic intoxication by inhalation of phosphorus fumes occasionally occurs in industry. A chronic phosphorus poisoning literature review for the period up to 1945 (and a presentation of recent case reports) summarizes the chronic effects of elemental phosphorus on humans (Heimann, 1945)⁴.

Almost all knowledge concerning the toxicity of white phosphorus to aquatic species derives from studies initiated in the late 1960's in response to a massive fish kill caused by the wastewater discharged from a factory manufacturing elemental phosphorus in Newfoundland (Idler, 1969⁵; Malin, 1972⁶). Most of the data concern salt water organisms. A similar

catastrophic die-off of freshwater fish in Pickwick Reservoir in the early 1950's is believed to have resulted from an accidental discharge of phosphy water from holding ponds at the Muscle Shoals, Alabama, phosphorus manufacturing facility operated by the Tennessee Valley Authority^{7,8,9}. However, phosphorus poisoning was never demonstrated conclusively, and very little research resulted from this episode. Because phosphine, phosphite, hypophosphite and hypophosphate are likely products of the decomposition of white phosphorus in water, the relevant chemistry and toxicology of these compounds are covered in the present report.

Approach to Mammalian Toxicology. A survey was conducted of the chemical and biological literature on white phosphorus and on the related compounds. Sources for the literature search are set out in detail in Appendix A. Most of the references were obtained from Chemical Abstracts. Several machine searches were also made, but were not very helpful. In addition to the references obtained from the sources listed, a paper by Fleming, Miller, and Swayne (1942)¹⁰ on "Some recent observations on phosphorus toxicology" noted that their references had been selected from some 1000 reports on phosphorus toxicology as having special bearing on their reported findings; this bibliography (actually only 444 references) was obtained from the technical library of the Tennessee Valley Authority.

The papers obtained were grouped in the following categories for toxicological evaluation.

1. Acute toxicity studies
2. Chronic toxicity studies
3. Clinical studies (as they relate)
4. Biochemical studies
5. Analytical and related chemical studies.

Further references on the chemistry and mammalian toxicology of white phosphorus, of only peripheral interest, are listed in Appendix B.

Approach to Toxicity to Aquatic Organisms. The principal source of information on the aquatic toxicology of white phosphorus is contained in a collection of reprints from Dr. David Idler of the Memorial University of Newfoundland⁷. Additional information and literature sources were forthcoming in telephone consultations with personnel from Memorial

University^{11,12}, ERCO Industries, Ltd.^{13,14}, Monsanto Company¹⁵⁻¹⁸, TVA⁷⁻⁹, Fisheries Research Board of Canada²⁰⁻²², American Chemical Society²³, and Ash Stevens, Inc.²⁴.

The published literature was independently surveyed using *Chemical Abstracts*. Appropriate subheadings under *phosphorus* were scanned from 1937 to August 1973. *Phosphine*, *phosphite*, *hypophosphite*, *hypophosphate*, *phosphorous acid*, *hypophosphorous acid*, and *hypophosphoric acid* were reviewed from 1917 to August 1973.

A machine search for relevant DOD documents was carried out using key words *phosphorus*, *white phosphorus*, *phossy water*, *plasticized white phosphorus*²⁵. Two documents relating to spray treatment of Pine Bluff phossy water, revealed as a result of this search, had not been received at the time this report was compiled. See also Appendix A.

Current Research on WP Toxicology and "Phossy Water". Apart from a few recent studies on the biochemical effects of white phosphorus, there does not appear to be any current research activity on the toxicity and toxicology of WP and phossy water. Laboratories listed in references 11, 12, 20, 21, and 22 are believed to be still active in research on the effects of elemental phosphorus pollution.

MAMMALIAN TOXICOLOGY OF WHITE PHOSPHORUS AND RELATED MATERIALS

Acute Toxicity in Animals. A summary of acute lethal toxicity of phosphorus in the rabbit, dog, rat, and mouse is given in Table 1. No LD50 values have been recorded for phosphorus. The effects of single toxic doses of phosphorus in olive oil given subcutaneously to male rabbits have been studied by Huruya (1928)²⁶. Changes associated with the production of fatty livers by white phosphorus in rats have been investigated by Seakins and Robinson (1964)²⁷ and by Pani, Gravela, Mazzarino and Burdino (1972)²⁸. Love (1934)²⁹ studied the effect of phosphorus on the normal and on the restored liver following partial hepatectomy in the albino rat.

Acute Toxicity in Man. In acute phosphorus poisoning, the predominant effect is fatty degeneration of the tissues. This occurs essentially in the liver, the damage to which is the cause of death in most cases. As little as 1/8 grain (8.1 mg) has been reported as having caused acute toxicity and death but the average fatal dose taken by mouth is generally considered to be of the order of 1.5 grains (97 mg). Christensen (1972)³⁷ gives the oral LD as 1.4 mg/kg. Acute phosphorus poisoning is peculiar because symptoms appear in two stages. During the first 24 hrs, symptoms

TABLE 1. Summary of Acute Toxicity of White Phosphorus in Animals.

Animal Species	Route of administration	Dose* (mg/kg)	Remarks	References
Rabbit	or	7	LD	Hirz (1913) ³⁰
	or	10	LD	Frank and Isaac (1910-1911) ³¹
	sc	12.5	LD	Santesson and Malmgren (1904) ³²
	sc	30	LD	Santesson and Malmgren (1904) ³²
	or	20.0	LD	Cameron and Patrick (1966) ³³
Dog	sc	2-3	LD	Rubow (1905) ³⁴
	sc	12	LD	Welsch (1905) ³⁵
Rat	or	<0.1 grain [†]	LD	Shepard (1951) ³⁶
	or	0.5	LD	Cameron and Patrick (1966) ³³
Mouse	or	3.5	LD	Cameron and Patrick (1966) ³³

*All administered in oil.

†For a 400 g rat.

of severe gastrointestinal irritation occur as soon as one-half hour after ingestion. The victim may die of cardiovascular failure within 12 hr. This first stage may be followed by a latent period lasting from a few hrs to a few days depending upon the amount ingested. The systemic stage is characterized by abdominal pain, nausea, vomiting, hematemesis and other hemorrhagic manifestations, jaundice, hepatomegaly, oliguria, toxic psychosis, convulsions, coma, and shock. There may be severe damage to the liver, heart, and kidney, and death may ensue at any time. Cirrhosis of the liver has been reported after recovery from the acute state. The signs and symptoms of acute poisoning in man are also adequately summarized by Gleeson, Gosselin, Hodge and Smith (1969)³⁸.

Fourteen papers on acute human phosphorus poisoning (episodes and case studies) are listed in the references (nos. 39-52) at the end of this report.

Chronic Toxicity in Animals. Sollmann (1925)⁵³ was the first investigator to study the effects of chronic feeding of phosphorus to albino rats. Further similar experiments by Flemming *et al.* (1942)¹⁰ confirmed the retardation in growth and the effects on bone development.

No significant changes were found in the blood, urine and tissues of dogs given subcutaneous injections of phosphorus in peanut oil (0.4, 0.2, and 0.1 mg P/kg/day) according to Buchanan, Sigal, Robinson, Pope, Ferguson and Thomison (1954)⁵⁴. The fate of orally administered choline in dogs with livers damaged through phosphorus poisoning was studied by Sigal, Buchanan and Robinson (1954)⁵⁵.

Ferraro, Jervis and English (1938)⁵⁶ drew attention to the fact that studies on white phosphorus intoxication in rabbits have been focused generally on liver changes and the metabolism of fatty substances. These workers have described the pathological changes in the brains of adult rabbits (nerve cells, blood vessels and glia) given iv injections of a 1% solution of phosphorus in oil (0.20 to 1.0 ml doses), over periods of 2 days to 15 weeks.

Lawrence and Huffman (1929)⁵⁷ noted that sc injections of phosphorus in olive oil into guinea pigs caused a marked increase in the number of the circulating monocytes in the blood.

Chronic Toxicity in Man (Including Potential for Carcinogenicity, Mutagenicity, and Teratogenicity). Chronic poisoning, seen mostly in factory workers exposed repeatedly to phosphorus vapors, is manifest

by cachexia, anemia, bronchitis, and necrosis of the mandible ("phossy jaw"). The signs and symptoms of chronic poisoning, whether from ingestion or inhalation, have been well documented in the publications of Heilmann (1946)⁴ and Gleeson, Gosselein, Hodge, and Smith (1969)³⁸. For summary of chronic toxicity to humans see Table 2.

The TLV for white phosphorus (ACGIH, 1972)⁵⁸ is 0.1 mg/m³. The TLV for phosphine⁵⁸ is 0.3 ppm or 0.4 mg/m³.

Testing for the possible carcinogenic activity of elemental phosphorus in experimental animals has been summarized by Hartwell (1951)⁵⁹, and Shubik and Hartwell (1957⁶⁰, 1969⁶¹). No animals with tumors were found following either oral feeding or subcutaneous injections in oil, even in rats receiving the compound for 33 months.

White phosphorus does not appear to have been investigated for possible mutagenic or teratogenic potential in living systems.

A case of ophthalmoplegia due to phosphorus poisoning has been described by Medea (1905)⁶².

The effect of white phosphorus on growing bones and growing teeth of both rabbits and rats has been studied by Adams and Sarrazat (1940)⁶³.

Burns due to Phosphorus. Two recent papers describe observations on the burns resulting from elemental white phosphorus. Curreri, Morris, and Pruitt (1970)⁶⁴ reviewed the clinical records of 111 patients, mostly with burns due to white phosphorus, who were admitted to the US Army Institute of Surgical Research. They noted the high incidence of residual ocular damage and suggested a number of specialized requirements for successful systemic and local wound care. No biochemical or toxicological data were evaluated.

Bowen, Whelan, and Nelson (1971)⁶⁵ describe a methodology for the production of a standard white phosphorus burn in rabbits. Although their study offered no suggestions regarding new or improved therapy of the white phosphorus-burned patient, it did elucidate electrolyte aberrations (depression of serum calcium and elevation of serum phosphorus and sometimes a reversal of the calcium/phosphorus ratio), and electrocardiographic changes which may explain early, sudden demise in the seemingly inconsequentially white phosphorus-burned patient.

Toxicity of Phosphine. Because of the possibility that phosphine (PH₃) may be a product of phosphorus metabolism (see p. 15) the toxicology has

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

phosphorus appears to be noncarcinogenic when fed to experimental animals. Testing for mutagenicity and teratogenicity has not been carried out. Many studies of a biochemical nature have been reported and have been summarized in the report. The primary effect of phosphine is neurological, and in this it differs markedly from white phosphorus.

White phosphorus is also highly toxic to aquatic animals. The 96 hr LC50's are less than 50 ppb for all fish studied, and the incipient lethal level is probably less than 1 ppb for most fish. Crustaceans and many molluscs are more tolerant, but still succumb to phosphorus concentrations of 1 ppm or less. Phosphorus poisoning appears to be cumulative and irreversible for fish and lobsters, though the cause of mortality in fish has not been determined. There is no information on the toxicity of phosphine to aquatic biota or experimental animals.

Recommendations for studies on white phosphorus (and phosphy water) in experimental animals and in wastewaters are outlined.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

TABLE 2. Summary of Chronic Toxicity of White Phosphorus to Humans
(Helmann, 1946)⁴.

System Affected	Characteristic Effects
Osseous	Jaw necrosis ("Phossy Jaw"). Hyperostosis of entire Skeletal Structure. Ossification of growing centers of bones.
Blood	Low in potassium and inorganic P content, high in chlorides, fat content. Morphological changes in circulating blood cells. Anemia of so-called "secondary" type.
Urinary	General cellular damage. Increase in ammonia N ₂ and unoxidized sulfur content.
Liver	No definite liver diseases described.
Local Irritant Effects	Skin burns, conjunctivitis, skin rashes.
General Cachexia	Exhaustion; loss of appetite; indefinite digestive complaints; garlic-like breath; slight jaundice, bleeding of the mucous membranes and albumin in the urine.

been summarized herewith (see also Webster, 1946⁶⁶). The toxicology of acute mammalian poisoning by the gas phosphine has not been fully worked out according to Patty, Fassett, and Irish (1963)⁶⁷. It does not alter the blood and its effects are apparently exerted largely through the central nervous system. Some inflammation of the lungs results from its inhalation. The acute action of phosphine does not resemble that of phosphorus; instead, there is marked dyspnea, purgation, weakness, tremor, and finally violent convulsions and death. Mild cases recover without after-effects. Prolonged exposure to small amounts of phosphine is said to give rise to chronic symptoms similar to those of phosphorus poisoning and may be explained by the decomposition of the compound in the body (Meissner, 1924⁶⁸; Müller, 1940⁶⁹; Loewenthal, 1949⁷⁰). A summary of the physiological responses to various concentrations of phosphine is given in Table 3.

BIOCHEMICAL STUDIES

Poisoning due to phosphorus, according to Cutler (1931)⁷², results in an increase of guanidine in the blood of dogs and accompanying hypoglycemia. The results also indicate that the guanidine increase is probably dependent upon liver damage rather than upon kidney failure.

Mizuno (1932)⁷³ reported a relatively large increase (3 to 8 times) in the lactic acid content of the urine of rabbits with liver damage due to phosphorus poisoning.

The glucuronic acid produced by surviving liver slices from guinea pigs given a sc injection of yellow phosphorus (0.75 mg/kg body wt) in olive oil was found by Bueding and Ladewig (1939)⁷⁴ to be considerably reduced. Chloroform poisoning did not affect glucuronic acid production.

Barone, Cittadini, Galeatti, and Terranova (1973)⁷⁵ reported that fatty liver degeneration in rats caused by white phosphorus had no influence on the level of the microsomal respiratory pigments cytochrome b₅ and cytochrome P₄₅₀.

Rabbits with fatty livers caused by white phosphorus poisoning had three times the amounts of cytochrome c over fasted control animals according to Prader (1947)⁷⁶. Milder poisoning resulted in less cytochrome c and a lag period was also evident.

Glosa⁵, Porta, and Hartroft (1972)⁷⁷ measured the liver microsomal glucose-6-phosphatase in the fatty livers of rats given oral doses of

TABLE 3. Summary of the Physiological Responses to Various Concentrations of Phosphine (Henderson and Haggard, 1943)⁷¹.

Response	Phosphine Concentration	
	ppm	ug/m ³
MAC for prolonged exposure	2	2.6
Slight symptoms after several hrs. exposure	7	10,010
Max. concentration that can be inhaled for 1 hr. without serious consequences	100 - 200	143,000 - 286,000
Dangerous after-exposure for 1/2 - 1 hr.	400 - 600	572,000 - 878,000
Fatal after-exposure for 30 mins.	1,000 - 2,000	1,430,000 - 2,860,000

phosphorus (7.5 mg/kg body wt) and found no significant depression at 6 or 24 hours.

Serological studies on the blood of rabbits given yellow phosphorus (0.1, 1.0, 2.0 mg/kg body wt) were reported by Tanaka (1960)⁷⁸ with the following results: All groups showed an early increase in non-protein nitrogen, amino acid nitrogen and uric acid, and a slight increase in creatine and creatinine. A decrease in erythrocyte count, hemoglobin and red cell fragility was also recorded.

The changes in the serum proteins after blood and plasma losses have been studied in normal and phosphorus poisoned rabbits by Wenner (1949)⁷⁹. He found an increase of the α - and β -globulins after bleeding but no increase of the glucosamine content.

Fleming and Collings (1951)⁸⁰ studied the chylomicron count in the blood of rats given sc injections of white phosphorus (1.1 mg/kg body wt) in peanut oil. No significant changes occurred in fasting rats but significant and reproducible changes were noted 4 hours after resuming food intake.

Recent papers by Truhaut *et al.* (1969⁸¹, 1971⁸², 1972⁸³) describe the effects of both single sc doses and repeated sc doses of white phosphorus in the rat. The effects on the levels of liver phospholipids and triglycerides were reported.

TOXICITIES OF WHITE PHOSPHORUS TO AQUATIC ORGANISMS

Toxicity of Elemental Phosphorus in Water. In the earliest aquatic study of which we are aware, the tolerance of bluegills (*Lepomis macrochirus* Rafinesque) for white phosphorus was measured in freshwater at 26°C (Isom, 1960)⁸⁴. The author concluded that phosphorus was nontoxic to the limit of its solubility (ca. 3 mg/l at 15°C), but paradoxically, that much smaller quantities of colloidal phosphorus were highly toxic. The conclusion concerning colloid-free water is probably invalid, because the author did not have available an analytical procedure sufficiently precise to tell whether elemental phosphorus was present at all; however, the 96 hour median tolerance level of 45 μ g/l, colloidal or dissolved, may reasonably be taken as an upper limit (cf. Table 1). Curiously, a Russian report claims that the toxicity of different forms of elemental phosphorus to freshwater fishes decreases in the order emulsion > solution > suspension > colloid (Kresmov, 1970)⁸⁵.

The most complete studies on white phosphorus intoxication in a marine environment have been carried out in connection with operation of a plant producing elemental phosphorus in Newfoundland^{5,6}. This plant, constructed by ERCO Industries, Ltd. on Long Harbour, Placentia Bay, began discharging phosphy water in December, 1968. Shortly thereafter large numbers of fish, principally cod and herring, were found dead in Long Harbour and adjacent waters, and in May, 1969, the plant was shut down and part of Placentia Bay was closed to fishing. Subsequently, groups associated with the Fisheries Research Board of Canada have studied the toxicity of white phosphorus in seawater to various fish and crustaceans (Zitko *et al.*, 1970⁸⁶; Fletcher *et al.*, 1970⁸⁷; Fletcher and Hoyle, 1972⁸⁸). Their results are summarized in Table 4. For five genera and four families of fish in saltwater, 96 hour LD 50's are remarkably uniform, $3 \pm .7$ $\mu\text{g/l}$, although field experience indicates that cod are substantially more tolerant than herring (Fletcher, 1973)¹². Incipient lethal levels are very low, suggesting that phosphorus poisoning is accumulative in these fish. Two separate studies emphasize the importance of the time-dosage relationship (Zitko *et al.*, 1970⁸⁶; Fletcher and Hoyle⁸⁸); a short exposure to a relatively high concentration may be lethal in the same time period as continuous exposure to a low concentration. For crustaceans the tolerance to phosphorus may be several orders of magnitude higher, though the incipient lethal level for lobsters is still quite low in terms of likely pollution levels. The beach flea has the greatest tolerance for white phosphorus of any animal studied, and is the only one for which reversibility of intoxication has been clearly demonstrated.

Similar investigations of other phyla have not been reported. However, a study of the bottom of Long Harbour in the vicinity of the effluent outlet provided evidence of selective mortality among benthic organisms (Peer, 1972)⁹¹. Thus, e.g., mussels (*Modiolus modiolus*) and a burrowing sea anemone (*Edwardsia* sp.) were alive and apparently normal in a region from which scallops and sand dollars (*Echinarachnius parina*) had been nearly exterminated. In a recent study, Black (1973)²² has found no significant variation in annual growth rates for Long Harbour mussels (*Mytilus edulis*) during 1969, the year of significant phosphorus pollution, and preceding and succeeding years (Black, 1973)⁹². On the other hand, no periwinkles (*Littorina littorea*) old enough to have been set before 1970 were found, indicating that all were killed by the 1969 spill.

Accumulation of Phosphorus by Aquatic Organisms. In a 16 hour exposure, cod have been shown to concentrate phosphorus 1000 times in the liver and 10 to 1000 times in muscle tissue from water containing 20 to 80 $\mu\text{g/l}$ (Dyer *et al.*, 1970)⁹³. Concentration factors are roughly proportional to the lipid content of the tissue, and inversely related to the phosphorus content of the water. Other fish have not been studied in such

TABLE 4. Summary of Toxicity of White Phosphorus to Fish Species and Crustaceans.

Organism	Incipient Lethal Level mg/l	96 hr LC50 mg/l	Test	Water	Temp. °C	References
Bluegill (<i>Lepomis macrochirus</i>)		ca. 45	static	fresh	26	Isom (1960) ⁸⁴
Cod (<i>Gadus morhua</i>)	<1.9	2.5	cont.	salt	9-10	Fletcher and Hoyle (1972) ⁸⁸
Atlantic salmon (<i>Salmo salar</i>)	<.8	2.3	cont.	salt	10-13	Zitko et al. (1970) ⁸⁶ Fletcher and Hoyle (1972) ⁸⁸
Herring (<i>Clupea harengus</i>)	<2.5	3.7	static	salt	7-9	Zitko et al. (1970) ⁸⁶
Brook trout (<i>Salvelinus fontinalis</i>)	<.5	2.5	cont.	salt	7-9	Fletcher et al. (1970, 1971) ^{87,89}
Smelt (<i>Osmerus mordax</i>)		2.5	cont.	salt	7-9	Fletcher et al. (1970, 1971) ^{87,89}
Killifish (<i>Fundulus heteroclitus</i>)		>20	cont.	salt	7-9	Fletcher et al. (1971) ⁸⁹
Cunner (<i>Tautoglabrus adspersus</i>)		*	cont.	salt	7-9	Fletcher et al. (1971) ⁸⁹
Lobster (<i>Homarus americanus</i>)	20-40	300	static	salt	?	Zitko et al. (1970) ⁸⁶ Fletcher (1971) ⁹⁰
Beach flea (<i>Gammarus oceanicus</i>)	3000-4000		static	salt	18-21	Zitko et al. (1970) ⁸⁶
Atlantic salmon smolts		ca. 10	cont.	fresh	11-14	Fletcher and Hoyle (1972) ⁸⁸




*LT 50, 3-4 µg/l, 200-300 hr.

detail, but from a survey of fish collected in Placentia Bay in 1969-1970, and from limited bioassays, it is known that herring and salmon accumulate phosphorus to the 1 ppm level, and that other fish can accumulate at least measureable quantities (Ackman *et al.*, 1970)⁹⁴.

Fletcher (1971)⁹⁰ has recorded the 48 hour uptake of white phosphorus by several marine invertebrates and seaweed from seawater containing $15 \pm 9 \mu\text{g/l}$ of white phosphorus (cf. Table 5). It may be noted that periwinkles, which are foraging grazers, accumulated greater amounts of phosphorus than the filter feeding molluscs. Fletcher (1971)⁹⁰ suggests that the former may ingest phosphorus with the seaweed. Intoxication was not reported for any of these invertebrates, and elemental phosphorus was eliminated from all the organisms of Table 6 within seven days following transfer to uncontaminated seawater.

Toxicity to Waterfowl. Dabbling ducks of the genus *Anas* feeding in areas of phosphy water discharge have been found to ingest lethal quantities of elemental phosphorus (Coburn *et al.*, 1950)⁹⁵. In a series of experiments conducted with captive black ducks and mallards, chronic and acute responses varied widely among individuals, but a single dose of 3 mg/kg body weight was sufficient to kill all ducks studied in 6 to 33 hours.

Toxicity of the Total Waste. Analysis of a phosphy water sample from Pine Bluff AAP showed it to contain 242 mg/l of phosphate and 208 mg/l of lower phosphorus oxides, as phosphorus (Blumbergs *et al.*, 1973)^{24,96}. These might include hypophosphate, phosphite, and hypophosphite, the latter two being known products of nonaqueous oxidation of phosphorus (Van Wazer, 1958)⁹⁷. No one of these is reported to be very toxic.



Nitrogen fixation by *Azotobacter vinelandii* is completely inhibited by 300 mg/l of phosphite (Bulen and Frear, 1957)⁹⁸ as P, ca. 10^{-2}M), while some other bacteria (e.g. *Escheria coli* and certain *Pseudomonas*, *Agrobacter*, *Rhizobium* and *Agrobacterium* species) can utilize phosphite for growth at the same concentration (Casida, 1960)⁹⁹, and *Pseudomonas fluorescens*

TABLE 5. Phosphorus Accumulation by Seaweed and Invertebrates (Fletcher, 1971)9%.

Organism	White Phosphorus (mg/g)	Conc. Factor Approx.
Seaweed		
<i>Fucus vesiculosus</i>	.33	22
<i>Fucus distichus</i>	.33	22
Invertebrates		
Clam (<i>Mya arenaria</i>)	.34	23
Quahog (<i>Arctica islandica</i>)	.26	17
Mussel (<i>Mytilus edulis</i>)	.16	11
Periwinkle (<i>Littorina littorea</i>)	.64	43
Starfish (<i>Asterias vulgaris</i>)	.40	27

converts phosphite to phosphate in excess of that required for growth. Still other micro-organisms, including some yeasts, are unable to utilize phosphite, but are unaffected by it. Robertson and Boyer (1956)¹⁰⁰ report that rat weanlings fed inorganic phosphite at 250 mg/kg on alternate days show some restriction of growth, but 125 mg/kg is without effect. Table 6 summarizes enzyme inhibition data for phosphite and hypophosphite at the $10^{-2}M$ level. The effect is generally small. Van Wazer (1958)¹⁰¹ says that "the phosphites exhibit considerable toxicity, acting on the nerve centers and abdominal glands; he also says "there is no evidence that hypophosphite exerts the slightest effect on mammals, and it passes through the mammalian system unchanged"¹⁰², but he presents no data more recent than 1902. There appear to be no toxicity data of any kind for hypophosphate. At higher pH levels, phosphine is formed from elemental phosphorus, and a small amount may be present in phosphy water. Phosphine is a toxic gas (MPC 0.3 ppm for humans) with a moderately low water solubility. It is alleged to be toxic to rainbow trout in concentrations exceeding 3.6 ppm (Daudoroff and Katz, 1950)¹⁰⁶. A few papers report the generation of phosphine in natural waters by anaerobic sediments (Datsko, 1958¹⁰⁷; Luning and Brohm, 1932¹⁰⁸, 1934¹⁰⁹), and one implies that it is highly toxic to fish¹⁰⁷, but otherwise there does not appear to be a literature of aquatic toxicology. It is rapidly degraded by the microorganisms in wet soil (Hilton and Robison, 1972)¹¹⁰.

Polyphosphoric acid, if formed in phosphy water, would not be expected to present a problem. Although it is a sequestering agent, like NTA, and might enhance the toxicities of any heavy metals in the water, it is readily biodegradable to orthophosphoric acid (Metcalf, 1973¹⁵, Karl-kroupa *et al.*, 1957)¹¹¹.

The phosphy water discharged in the manufacture of phosphorus is a more complex waste than that released in filling operations, such as that at Pine Bluff AAP⁵,¹⁵⁻¹⁹. The latter consists only of dissolved and suspended elemental phosphorus and oxidation and hydrolysis products thereof (principally phosphoric acid). The manufacturing wastewater includes as well a large amount of suspended solids, fluosillicic acid and sulfur dioxide, and some ammonia and cyanide. A typical wastewater analysis for the ERCO Industries, Ltd. plant, prior to shutdown in 1969, is given in Table 7. Thus although extensive mortality studies were carried out in the area, it was necessary that early investigators consider that the catastrophic effect of the Long Harbour discharge may have been due in part to waste components other than phosphorus. Bioassays of the various waste constituents, however, have shown phosphorus to be the most toxic by several orders of magnitude (Fletcher *et al.*, 1971)⁸⁹. For example, Figure 1 illustrates the comparative toxicities of several substances to brook trout maintained in seawater. The total wastewater

TABLE 6. Enzyme Inhibition by Phosphite and Orthophosphite.

Enzyme	Anion	Conc	& Inhibition	References
Intestinal phosphatase	orthophosphite	$10^{-2}M$	5	Robertson and Boyer (1956) ¹⁰⁰
3-Phosphoglyceraldehyde dehydrogenase	orthophosphite	$2 \times 10^{-2}M$	salt effect only	Robertson and Boyer (1956) ¹⁰⁰
<i>E. coli</i> formic dehydrogenase	hypophosphite	$10^{-2}M$	~20	Crewther (1956) ¹⁰³
<i>E. coli</i> formic hydrogenlyase	hypophosphite	$10^{-2}M$	~40	Crewther (1956) ¹⁰³
<i>E. coli</i> hydrogenase	hypophosphite	?	0	Crewther (1956) ¹⁰³ Grubberg-Monago <i>et al.</i> (1951) ¹⁰⁴
Horse serum cholinesterase	hypophosphite	10^{-2}	~35	Frommel <i>et al.</i> (1944) ¹⁰⁵

TABLE 7. Analysis of Long Harbour Wastewater (Idler, 1969)⁵.

Flow	ca. 8000 gpm	11.5 MGD
Fluoride	22800 lb/day	245 mg/l
SO ₂	6250 lb/day	67 mg/l
Total P	1630 lb/day	18 mg/l
P ₄	1250 lb/day	13 mg/l
CN	96 lb/day	1 mg/l
NH ₃	455 lb/day	5 mg/l
Solids	38360 lb/day	410 mg/l




FIGURE 1: TOXICITIES OF VARIOUS SUBSTANCES TO SEAWATER MAINTAINED BROOK TROUT (Fletcher, *et al.*, 1971)⁸⁹

exhibits no enhanced toxicity to cod beyond that to be predicted on the basis of its white phosphorus content alone (Idler, 1965)⁵.

Other constituents of the waste may affect the distribution and dispersal of phosphorus (Jangaard, 1970)¹¹². Fluosilicic acid reacts with calcium and magnesium in seawater to produce a flocculant precipitate, which can coagulate and thereby immobilize colloidal white phosphorus in the immediate vicinity of the outfall. This greatly reduces the efficacy of tidal dilution, and may result in the creation of permanent toxic zones. In the case of the Long Harbour operation, much of the phosphorus contamination resulted from accidental spills in the course of loading ships with liquid phosphorus¹³. Phossey water from this source would not be expected to differ greatly from that from ammunition plants.

Behavioral Effects. The most obvious symptom of phossey water poisoning, other than death of the victim, is the red color acquired by the belly and head of certain fish, particularly herring. All fish eventually become lethargic, and on the approach of death may swim in circles (Zitko *et al.*, 1970)⁸⁶. Lobsters exposed to lethal levels of white phosphorus "lose muscle tone, become lethargic, move the pereopods in a jerky, uncoordinated and aimless manner, and eventually lose all neuromuscular response"⁸⁶. Beach fleas become paralyzed when subjected to lethal doses, but may recover when transferred to clean water⁸⁶.

Among waterfowl, acute poisoning results in depression, followed by leg weakness and finally violent convulsions and death. Birds suffering from chronic poisoning steadily lose weight and eventually show signs of paralysis. Field cases, usually suffering from acute poisoning, are described as having "ungainly flight" (Coburn *et al.*, 1950)⁹⁵.

Pathology. White phosphorus can be expected to cause extensive tissue damage wherever it accumulates. Tissues of the gills, kidneys, liver and spleen may undergo substantial disintegration (Odense *et al.*, 1972)¹¹³. However, the symptoms of phosphorus intoxication, including death, are not necessarily related to tissue destruction.

The cause of fish mortality from phossey water has yet to be determined. White phosphorus levels low enough to permit survival of the fish for more than a few hours may cause extensive hemolysis of erythrocytes for some species, viz. herring (Zitko *et al.*, 1970⁸⁶; Fletcher *et al.*, 1970⁸⁷), Atlantic salmon (Zitko *et al.*, 1970⁸⁶; Fletcher and Hoyle, 1972⁸⁸), and brook trout (Fletcher *et al.*, 1970⁸⁷). Some of the hemoglobin released may then leave the circulatory system and pass into the interstitial areas, temporarily imparting a bright red color to the head and belly. Hematocrits

may drop to a very low value, and in latter stages the gills may turn pale, but loss of red cells is apparently not the cause of death. Cod and smelt exhibit no evidence of hemolysis at any level of phosphorus exposure, and trout and salmon expire without hemolysis if the dosage is great enough (Fletcher *et al.*, 1970⁸⁷, 1972⁸⁸). Recent Russian studies suggest that the central nervous system may be involved (Krasnov, 1970⁸⁵; Mazmanidi, 1970¹¹⁴). Fletcher (1973)¹¹⁵ has found that brook trout fed cod poisoned by elemental phosphorus develop the same symptoms as trout exposed to phosphorus in water. Whatever the cause, phosphorus intoxication of fish appears not to be reversible.

In lobsters, phosphorus intoxication brings about degeneration of the hepatopancreas and antennal gland tissue (Aiken and Byard, 1972)¹¹⁶. Although this damage should eventually kill the lobster, death is probably due to asphyxiation resulting from blood coagulation. Toxic effects are cumulative and irreversible or very slowly reversible.

For waterfowl, the histopathology of phosphorus intoxication is inconsistent. The kidneys are usually congested, the liver commonly shows signs of damage, and duodenal ulcers may be present, but it is not certain that any of these disorders are related to the early death of the bird (Coburn *et al.*, 1950)⁹⁵.

Summary and Conclusions. White phosphorus is highly toxic to fish. The 96 hour LC 50's are less than 50 ppb for all fish studied, and the incipient lethal level is probably less than 1 ppb for most fish. Phosphorus poisoning appears to be accumulative and irreversible, though the cause of mortality has not been determined. While phosphorus is readily taken up by fish and other aquatic organisms directly from the water, fish may also acquire lethal quantities of elemental phosphorus through the food chain, since the few macroinvertebrates studied have a much higher tolerance for white phosphorus than fish. The symptoms of phosphorus intoxication are passed onto brook trout when they are fed muscle tissue from phosphorus-poisoned cod (Fletcher, 1973)¹¹⁵. Furthermore, elemental phosphorus can be passed on to humans, since a considerable portion, 25% or more, remains in the muscle of the fish after processing, storage, and cooking (Dyer *et al.*, 1972)¹¹⁷.

So toxic is white phosphorus to some fish that it is difficult to set maximum safe levels, although there are indications that fish acquire a tolerance after exposure (Fletcher, 1973)¹². If it is assumed that the minimum incipient lethal level is 0.1 µg/l (1/5 the lowest measured), then it would seem advisable to set a maximum of 0.01 µg/l in the mixing zone of any outfall. Zero discharge of phosphorus manufacturing process water has been recommended (Anon., 1973)¹¹⁸.

Other components of phosphy water are probably inconsequential in terms of their likely concentrations, but there is uncertainty concerning phosphine.

CHEMISTRY OF WHITE PHOSPHORUS AND DERIVED SUBSTANCES

The Fate of Elemental Phosphorus in Water. Although there are no definitive published reports on the kinetics of oxidation of elemental phosphorus in water, it appears that the rate is highly dependent on the degree of dispersion. At concentrations (ca. 10 µg/l) well below the accepted solubility limit of 3 mg/l, with the dissolved oxygen content unspecified, elemental phosphorus disappears by a first order process with a half-life of 2 hours at ca. 10°C (Zitke *et al.*, 1970)⁸⁶ 0.85 hour at 30°C (Addison, 1971)¹¹⁹. At concentrations (50 to 100 mg/l) well above the solubility limit, with a dissolved oxygen content of 6 to 7 mg/l, the same reaction has a half-life of 80 hours at 30°C and 240 hours at 0°C (Bullock and Newlands, 1969)¹²⁰. The relatively small temperature effect combined with the large inverse concentration effect is consistent with a diffusion controlled process. The oxidation of colloidal phosphorus in seawater is reported to be measurably slower than in fresh water, suggesting that the high salt content brings about agglomeration of the phosphorus particles¹²⁰. Thus, rapidly moving fresh water should lose elemental phosphorus faster than quiescent seawater.

Addison has presented evidence indicating that 1 ppm aqueous elemental phosphorus is oxidized to phosphate in a single step¹¹⁹. Likely intermediates, though stable under the reaction conditions, could not be isolated. Blumbergs *et al.*, 1973^{24,96} on the other hand, report that phosphy water from Pine Bluff AAP contains as much phosphorus in intermediate oxidation states as orthophosphate. A synthetic mixture prepared with several hundred ppm white phosphorus gave the same results. It is possible that the products of oxidation of suspended phosphorus are different from those of dissolved phosphorus.

The rapid reaction of white phosphorus with chlorine in water to form phosphorus acid *via* phosphorus trichloride is well documented (Blumbergs *et al.*, 1973⁹⁶; Anon, 1952¹²¹; Barber, 1969¹²²).

ANALYTICAL METHODOLOGY

Elemental Phosphorus. Until recently, the usual method for determining elemental phosphorus involved extraction of the sample with an organic

solvent such as benzene, oxidation of the extracted phosphorus to orthophosphate, and estimation of the latter by the molybdenum blue or other colorimetric method. Under ideal conditions, a detection limit of about 0.01 ppm of phosphorus in water is attainable by this technique, but of course it is not possible to distinguish P_4 from benzene-soluble phosphorus derivatives, phosphine in particular (McGilvery, 1973)¹⁴.

Addison and Ackman have devised a gas-liquid chromatographic procedure for elemental phosphorus capable of detecting as little as $10^{-12}g$ (Addison, 1971¹¹⁹; Ackman and Addison, 1969¹²³; Addison and Ackman, 1970¹²⁴). Benzene extracts of mud, water, or biological samples are injected onto a gas-liquid chromatograph equipped with a flame photometric detector. Techniques have been contrived to permit detection of 0.01 $\mu g/l$ of elemental phosphorus in water. (The solubility of P_4 in benzene is 10^4 times its solubility in water). Some interference from organics has been noted (Addison, 1971)¹¹⁹; for natural water samples, $\pm 0.1 \mu g/l$ is probably the limit of precision, and for mud and biological samples, $\pm 0.1-1 mg/kg$; depending on the nature of the sample. Using their methods, the authors and associates have analyzed for phosphorus in water (Addison *et al.*, 1971¹²⁵, 1972¹²⁶), mud (Ackman *et al.*, 1972^{126,127}), and fish (Ackman *et al.*, 1970^{94,128}) samples from Long Harbour.

More recently, Bohl and Kaelble (1972)¹²⁸ of Monsanto have devised a method for the determination of white phosphorus in air, also employing flame photometric gas chromatography. They cite a detection limit of $10^{-11}g$ for P_4 dissolved in xylene. Gorzny (1972)¹²⁹ has reviewed methods for elemental phosphorus through early 1970. It appears at this time that flame photometric gas chromatography is the method of choice for environmental samples.

Phosphine. Methods for phosphine have been reviewed by Fetchin and Grayson (1972)¹³⁰. Because phosphine is usually encountered as a gas, analytical procedures are based on air sampling. A typical procedure utilizes formation of a colored complex with silver diethyldithiocarbamate (Dechant *et al.*, 1966)¹³¹, the sensitivity corresponding to a level of ± 50 ppm in water. Berck *et al.*, 1970¹³² employing flame photometric gas chromatography, are able to detect quantities of gaseous phosphine in the picogram ($10^{-12}g$) range. Extraction and gas chromatography should be applicable to water samples containing phosphine.

Oxyacids of Phosphorus. Van Wazer (1958)¹³³ and Ohashi (1972)¹³⁴ have reviewed analytical methods for the oxyacids of phosphorus. The preferred method is separation by paper on thin layer (Seiler, 1965)¹³⁵ chromatography and estimation of the developed spots. More recently, Addison

has employed a gas chromatographic technique based on formation of trimethylsilyl derivatives and utilizing the flame ionization detector (Addison, 1971¹¹⁹; Butts, 1970¹³⁶). Sensitivities in the 0.1 ppm range are implied.

RECOMMENDATIONS FOR FUTURE EXPERIMENTAL AND FIELD STUDIES ON WHITE PHOSPHORUS

1. The determination of the LD50 for white phosphorus should be undertaken in two animal species.
2. Because there is no information on the long-term effects of phosphorus, both chronic 90-day studies of low dose levels of phosphorus in 2 species of animals as well as a 2 yr long-term feeding study of very low dose levels of phosphorus in animal species should be undertaken, with special attention being paid to bone development.
3. The metabolism of phosphorus using radio active ^{32}P in animal species should be studied especially to determine retention in bone and possibly in other organs. The form of urinary and fecal excretion of phosphorus should be investigated.
4. Teratogenic (especially for possible interference with bone development) and mutagenic studies of white phosphorus might be desirable.
5. An investigation of phosphine (PH_3) production from elemental phosphorus by microorganisms in both field and laboratory studies should be undertaken.
6. The recommendation made by Rosenblatt, Small, and Barkley (1973)¹ in their report (p. 25) viz. "Procedures for monitoring white phosphorus in water should continue to be developed. The toxicity and analysis of lower oxides of phosphorus and polyphosphates should be investigated", is fully supported in this report.
7. Because elemental phosphorus is by far the most toxic constituent, the primary data required for the assessment of phosphy water pollution is the rate of oxidation of dissolved phosphorus in the concentration range of 1-1000 $\mu\text{g/l}$. A definitive kinetic study should be initiated to determine the effects of phosphorus concentration, dissolved oxygen content, temperature, pH and dissolved solids. From such data the time required to achieve safe phosphorus levels could be estimated for any particular body of water.

8. Pine Bluff AAP phosphy water should be analyzed for the presence of intermediate oxyacids of phosphorus and for phosphine. Acute toxicities, based on LD 50's, should be measured for those present at the mg/l level, using appropriate freshwater target species, such as fish and crayfish. For any substance found to be toxic in terms of the quantities present in the waste microbiological studies should be initiated to measure inhibition and utilization of phosphorus.

9. Although toxicity data for elemental phosphorus is available for numerous species of sea-water fish and a few lower animals, further bioassay studies on phosphy water using freshwater fish, lower animals and algae is desirable. Chronic toxicity data for white phosphorus levels below 0.1 ppb would be useful for establishing maximum safe concentrations, but the rapid oxidation of dissolved phosphorus at these levels and uncertainties in the present best analytical method would appear to present awesome experimental complications.

LITERATURE CITED

1. Rosenblatt, D. H., M. J. Small, and J. J. Barkley, "Munitions Production Products of Potential Concern as Waterborne Pollutants Phase 1", US Army Medical Environmental Engineering Research Unit report No. 73-07, June 1973, pp. 23-31.
2. Oliver, T., "The Conquest of An Occupational Disease: Phosphorus Poisoning in Lucifer Match Making", *J. Roy. Inst. Pub. Hlth. Hyg.*, 1:660-666 (1938).
3. Hamilton, A. and H. L. Hardy, *Industrial Toxicology*, Second Edition, pp. 138-144, P. B. Hoeber, Inc., New York, 1949.
4. Heimann, H., "Chronic Phosphorus Poisoning", *J. Ind. Hyg. Toxicol.*, 28:142-150 (1946).
5. Idler, D. R., "Coexistence of a Fishery and a Major Industry in Placentia Bay", *Chem. Can.*, 21:16-21 (1969).
6. Malin, H. M., Jr., "Waste Control in a Fragile Environment", *Environ. Sci. Tech.*, 6:980-982 (1972).
7. Nicholas, W. R., Water Quality Branch, Div. of Environmental Planning, TVA, Chattanooga, TN (615) 755-3011. Personal communication, July, 1973.
8. Chance, C. Jack, Chief of Fisheries, TVA, Norris, TN (615) 494-7173. Personal communication, July, 1973.
9. Isom, Billy G., Environmental Biology Branch, Division of Environmental Research and Development, TVA, Muscle Shoals, AL (205) 383-4631. Personal communication, July, 1973.
10. Fleming, R. B. L., J. W. Miller, and V. R. Swayne, Jr., "Some Reagent Observations on Phosphorus Toxicology", *J. Ind. Hyg. Toxicol.*, 24:154-158 (1942).
11. Idler, Dr. David R., Marine Science Laboratory, Memorial University of Newfoundland, St. Johns, Newfoundland, Canada (709) 726-6681. Personal communication, July, 1973.
12. Fletcher, Dr. Garth L., Marine Science Laboratory, Memorial University of Newfoundland, St. Johns, Newfoundland, Canada (709) 726-6681. Personal communication, July, 1973.

LITERATURE CITED (Cont)

13. James, K. M., Manager, Marketing Services, ERCO Industries Ltd., Islington, Ontario, Canada (416) 239-7111. Personal communication, July, 1973.
14. McGilvery, Dr. James D., Manager, Research Department, ERCO Industries Ltd., Islington, Ontario, Canada (416) 239-7111. Personal communication, July, 1973.
15. Metcalf, Joe, Monsanto Co., St. Louis, MO (314) 694-2024. Personal communication, July, 1973.
16. Mullins, Morton, Monsanto Co., St. Louis, MO (314) 694-2648. Personal communication, July, 1973.
17. Jewell, John P., Supervisor, Pollution Control, Monsanto Co., Columbia, TN (615) 388-3431. Personal communication, July, 1973.
18. Bohl, Dr. Carl D., Monsanto Co., St. Louis, MO (314) 694-2648. Personal communication, July, 1973.
19. Derr, Dr. Paul F., Asst. Director, Research and Development, FMC Corp., Princeton, NJ (609) 452-2300. Personal communication, July, 1973.
20. Ackman, Dr. R. G., Fisheries Research Board of Canada, Halifax Laboratory, Halifax, Nova Scotia, Canada (902) 423-1293. Personal communication, July, 1973.
21. Addison, Dr. R. F., Marine Ecology Laboratory, Bedford Institute, Dartmouth, Nova Scotia, Canada (902) 426-3279. Personal communication, July, 1973.
22. Black, Dr. Robert, Fisheries Research Board of Canada, Biological Station, St. Johns, Newfoundland, Canada (709) 726-5330. Personal communication, July, 1973.
23. Malin, H. Martin, Jr., Assoc. Editor, Environmental Science and Technology, American Chemical Society, Washington, DC (202) 872-4586. Personal communication, July, 1973.
24. Blumbergs, Dr. P., Ash Stevens Inc., Detroit, MI (313) 872-6400. Personal communication, July, 1973.
25. Defense Documentation Center, Defense Supply Agency. Cameron Agency, Alexandria, VA 22314.

LITERATURE CITED (Cont)

26. Huruya, A., "Studies on the Phosphorus Intoxication of Rabbits", *J. Biochem. (Japan)* 10:63-95 (1928).
27. Seakins, A. and D. S. Robinson, "Changes Associated with the Production of Fatty Livers by White Phosphorus and by Ethanol in the Rat", *Biochem. J.*, 92:308-312 (1964).
28. Pani, P., E. Gravela, C. Mazzarino, and E. Burdino, "On the Mechanism of Fatty Liver in White Phosphorus-Poisoned Rats", *Exp. Mol. Pathol.*, 16:201-209 (1972).
29. Love, J. G., "Experimental Pathology of the Liver. XI. The Effect of Phosphorus on the Normal and on the Restored Liver Following Partial Hepatectomy in the Albino Rat", *Arch. Pathol.*, 14:637-648 (1932).
30. Hirz, O., "Über den Einfluss des Phosphors auf den respiratorischen Stoffwechsel", *Z. Biol.*, 60:187-310 (1913).
31. Frank, E. and S. Isaac, "Über das Wesen des gestörten Stoffwechsels bei der Phosphorvergiftung", *Arch. Exp. Pathol. Pharmacol.*, 64:274-292 (1910-1911).
32. Santesson, C. G. and R. I. Malmgren, "Über die Wirkung des Phosphoresquisulfides", *Skand. Arch. Physiol.*, 15:229-237 (1904).
33. Cameron, J. M. and R. S. Patrick, "Acute Phosphorus Poisoning - The Distribution of Toxic Doses of Yellow Phosphorus in the Tissues of Experimental Animals", *Med. Sci. Law*, 8:209-214 (1966).
34. Rubow, V., "Über den Lecithingehalt des Herzens und der Nieren unter normalen Verhältnissen, un Hungerzustande und bei der fettigen Degeneration", *Arch. Exp. Pathol. Pharmacol.*, 52:173-204 (1905).
35. Welsch, H., "Research on the Pathogenesis of the Anatomical Lesions in Acute Phosphorus Poisoning", *Arch. Inter. Pharmacodyn.*, 14:211-265 (1905).
36. Shepard, H. H., "The Chemistry and Action of Insecticides", p. 102 McGraw Hill Company, New York (1951).
37. Christensen, H. E. (ed.), "Toxic Substances List", 1972 ed., p. 422, USAHEW/NIOSH (1972).

LITERATURE CITED (Cont)

38. Gleason, M. N., R. E. Gosselin, H. C. Hodge, and R. P. Smith, "Clinical Toxicology of Commercial Products. Acute Poisoning", Third Edition, pp. 192-195, The Williams & Wilkins Co., Baltimore (1969).
39. Hann, R. G. and R. A. Veale, "A Fatal Case of Poisoning by Phosphorus, with Unusual Subcutaneous Hemorrhages", *Lancet*, 1:163-164 (1910).
40. Tidy, H. L., "The Relation of Acute Phosphorus Poisoning to Acidosis", *Lancet*, 1:19 (1911).
41. Dwyer, H. L., "Phosphorus Poisoning in a Child from the Ingestion of Fireworks", *J. Amer. Med. Assoc.*, 84:1254-1256 (1925).
42. Swinton, C. F., "Acute Phosphorus Poisoning", *Brit. Med. J.*, 2:1080 (1927).
43. Rossier, P. H., "Contribution à l'étude de l'intoxication aigue par le Phosphore", *Schweiz. Med. Wochensch.*, 57:890 (1927).
44. Blumenthal, S. and A. Lesser, "Acute Phosphorus Poisoning", *Amer. J. Diseases Children*, 55:1280-1287 (1938).
45. Chretien, T. E., "Acute Phosphorus Poisoning - Report of a Case with Recovery", *New Engl. J. Med.*, 232:247-249 (1945).
46. Newburger, R. A., S. B. Beaser, and H. Shwachman, "Phosphorus Poisoning with Recovery Accompanied by Electrocardiographic Changes", *Amer. J. Med.*, 4:927-930 (1940).
47. Rubitsky, H. J. and R. M. Myerson, "Acute Phosphorus Poisoning:", *Arch. Int. Med.*, 83:164-178 (1949).
48. Díaz-Rivera, R. S., P. J. Collazo, E. R. Pons, and M. V. Torregrosa, "Acute Phosphorus Poisoning in Man: A Study of 56 Cases", *Medicine*, 29:269-298 (1950).
49. Wechsler, L. and R. L. Wechsler, "Phosphorus Poisoning: The Latent Period and Unusual Gastrointestinal Lesions", *Gastroenterology*, 17:279-283 (1951).
50. Caley, J. P. and I. A. Kellock, "Acute Yellow Phosphorus Poisoning With Recovery", *Lancet*, 1:539-541 (1955).

LITERATURE CITED (Cont)

51. Cameron, J. M. and E. Rentoul, "Acute Phosphours Poisoning", *Med. Sci. Law*, 3:71-76 (1963).
52. Talley, R. C., J. W. Linhart, A. J. Trevino, L. Moore, and B. M. Beller, "Acute Elemental Phosphorus Poisoning in Man: Cardio-vascular Toxicity", *Amer. Heart J.*, 84:139-140 (1972).
53. Sollmann, T., "Studies of Chronic Intoxications on Albino Rats. VIII. Yellow Phosphorus", *J. Pharmacol. Exp. Ther.*, 24:119-122 (1925).
54. Buchanan, D. J., M. V. Sigal, C. S. Robinson, K. Pope, M. E. Ferguson, and J. B. Thomison, "Studies of Phosphorus Intoxication. I. Changes in Blood, Urine, and Tissues of Dogs Poisoned with Phosphorus", *Arch. Ind. Hyg. Occupational Med.*, 9:1-8 (1954).
55. Sigal, M. V., D. J. Buchanan, and C. S. Robinson, "Studies on Phosphorus Intoxication. II. Fate of Orally Administered Choline in Dogs with Damaged Liver", *Arch. Ind. Hyg. Occupational Med.*, 9:142-147 (1954).
56. Ferraro, A., G. A. Jervis, and W. H. English, "Pathological Changes in the Brain in Cases of Experimental Phosphorus Intoxication", *Psychiat. Quart.*, 12:294-305 (1938).
57. Lawrence, J. S. and M. M. Huffman, "An Increase in the Number of Monocytes in the Blood Following Subcutaneous Administration of Yellow Phosphorus in Oil", *Arch. Pathol.*, 7:813-819 (1929).
58. American Conference of Governmental Industrial Hygienists, "Threshold Limit Values of Airborne Contaminants Adopted by ACGIH for 1972", ACGIH, Cincinnati, Ohio (1972).
59. Hartwell, J. L., "Survey of Compounds Which have been Tested for Circinogenic Activity", Second Edition, p. 12 (1951), PHS Pub. No. 149.
60. Shubik, P. and J. L. Hartwell, "Survey of Compounds Which have been Tested for Carcinogenic Activity", Suppl. 1, p. 21 (1957). PHS Pub. No. 149.
61. Shubik, P. and J. L. Hartwell, "Survey of Compounds Which have been Tested for Carcinogenic Activity", Suppl. 2, p. 20 (1969). PHS Pub. No. 149.

LITERATURE CITED (Cont)

62. Medea, E., "Un caso di oftalmoplegia da avvelenamento fosforico", *Riv. Sper. di Freniat. Riggio-Emilia*, 31:246-247 (1905).
63. Adams, C. O. and B. G. Sarnat, "Effects of Yellow Phosphorus and Arsenic Trioxide on Growing Bones and Growing Teeth", *Arch. Pathol.*, 30:1192-1202 (1940).
64. Curreri, P. W., J. Morris, and B. A. Pruitt, Jr., "The Treatment of Chemical Burns: Specialized Diagnostic, Therapeutic, and Prognostic Considerations", *J. Trauma*, 10:634-642 (1970).
65. Boven, T. E., T. J. Whelan, Jr., and T. G. Nelson, "Sudden Death After Phosphorus Burns: Experimental Observations of Hypocalcemia, Hyperphosphatemia and Electrocardiographic Abnormalities Following Production of a Standard White Phosphorus Burn", *Ann. Surgery*, 174:779-784 (1971).
66. Webster, S. H., "Volatile Hydrides of Toxicological Importance", *J. Ind. Hyg. Toxicol.*, 28:167-182 (1946).
67. Patty, F. A., D. W. Fassett, and I. D. Irish (Eds.), "Industrial Hygiene and Toxicology, Vol. 2", pp.883, *Toxicology*, Interscience Publishers, New York, 1963.
68. Meissner, R., "Zur Toxikologie des Phosphorwasserstoffes", *Z. ges. Exptl. Med.*, 42:267-274 (1924).
69. Müller, W., "Über Phosphorwasserstoffvergiftung (Tierversuche) I. Mitteilung: Akute and Subacute Vergiftung", *Arch. Exp. Path. Pharmacol.*, 195:184-193 (1940).
70. Loewenthal, M., "Phosphine Poisoning", *Schweiz. Z. Path. u. Bakt.*, 12:313-350 (1949).
71. Henderson, Y. and H. W. Haggard, "Noxious Gases (and the Principles of Respiration Influencing Their Action)", Second Edition, p. 243, Reinhold Publishing Co., New York, 1943.
72. Cutler, J. T., "The Accumulation of Guanidine in the Blood Following Acute Liver Injury by Carbon Tetrachloride, Chloroform, Arsenic or Phosphorus", *J. Pharmacol. Exp. Ther.*, 41:337-345 (1931).

LITERATURE CITED (Cont)

73. Mizuno, H. "The Lactic Acid Content of the Urine of Patients with Liver Disease and of Rabbits with Experimentally Injured Liver", Japan. *J. Gastroenterol.*, 4:84-93 (1932). (C.A. 27:1046).
74. Bueding, E. and P. Ladewig, "Glucuronic Acid Produced by Surviving Slices of Liver From Animals Poisoned with Phosphorus or Chloroform", *Proc. Soc. Exp. Biol. Med.*, 42:464-465 (1939).
75. Barone, C., A. Cittadini, T. Galeotti, and T. Terranova, "The Effect of Intoxication Induced in Rat Liver by Carbon Tetrachloride, Ethionine and White Phosphorus on the Level of Microsomal Cytochromes b_5 and P_{450} ", *Experientia*, 29:73-74 (1973).
76. Prader, A., "Cytochrom-c-Gehalt der toxisch bedingten Fettleber", *Experientia*, 3:459-460 (1947).
77. Ghoshal, A. K., E. A. Porta, and W. S. Hartroft, "Comparative Studies on the Hepatotoxicity of CCl_4 and White Phosphorus in Rats", *Gastroenterol.*, 62:193 (1972).
78. Tanaka, N., "Serological Studies on Yellow Phosphorus Poisoning", (in Japanese) *Nichidai Igaku Zasshi*, 19:1537-1548 (1960).
79. Werner, I., "On the Regeneration of Serum Polysaccharide and Serum Proteins in Normal and Intoxicated Rabbits", *Acta Physiol. Scand.*, 19:27-42 (1949).
80. Fleming, R. B. L. and G. H. Collings, Jr., "The Chylomicron Count as an Indicator of Phosphorus Poisoning", *Arch. Ind. Hyg. Occup. Med.*, 4:567-572 (1951).
81. Truhaut, R., J. R. Claude, and J. M. Warnet, "Étude des modifications biochimiques provoquées par des intoxications aiguës et subaiguës au phosphore blanc chez le Rat. I. Action de doses répétées administrées par voie parentérale", *Ann. Pharm. Franç.*, 27:17-23 (1969).
82. Truhaut, R., J. R. Claude, and J. M. Warnet, "Étude des modifications biochimiques provoquées par des intoxications aiguës et subaiguës au phosphore blanc chez le Rat. II. Action d'une dose unique administrée par voie parentérale", *Ann. Pharm. Franç.*, 30:35-44 (1972).

LITERATURE CITED (Cont)

83. Truhaut, R., J. R. Claude, and J. M. Warnet, "Étude de l'incorporation de l'acide palmitique-³H-9-10 dans les triglycérides et les lipoprotéines du rat intoxiqué par le phosphore blanc", *C. R. Acad. Sci. Paris*, 272:2400-2403 (1971).
84. Isom, B. G., "Toxicity of Elemental Phosphorus", *J. Water Poll. Control. Fed.*, 32:1312-1316 (1960).
85. Krasnov, S. K., "Effect of Elementary Phosphorus on Fish (Pathophysiological Studies)", *Vop. Vod. Toksikol.*, 1970, 125; (C.A., 74, 122039t, 1971). Translation in prepn. by Nat. Res. Council, Canada.
86. Zitko, V., D. E. Aiken, S. N. Tibbo, K. W. T. Besch, and J. M. Anderson, "Toxicity of Yellow Phosphorus to Herring (*Clupea harengus*), Atlantic Salmon (*Salmo salar*), Lobster (*Homarus americanus*), and Beach Flea (*Gammarus oceanicus*)", *J. Fish. Res. Bd. Canada*, 27:21-29 (1970).
87. Fletcher, G. L., R. J. Hoyle, and D. A. Horne, "Yellow Phosphorus Pollution: Its Toxicity to Seawater - Maintained Brook Trout (*Salvelinus fontinalis*) and Smelt (*Osmerus mordax*)", *J. Fish. Res. Bd. Canada*, 27:1379-1384 (1970).
88. Fletcher, G. L. and R. J. Hoyle, "Acute Toxicity of Yellow Phosphorus to Atlantic Cod (*Gadus morhua*) and Atlantic Salmon (*Salmo salar*) Smolts", *J. Fish. Res. Bd. Canada*, 29:1295-1301 (1972).
89. Fletcher, G. L., R. J. Hoyle, and D. A. Horne, "The Relative Toxicities of Yellow Phosphorus Production Wastes to Seawater Maintained Fish", *Fish. Res. Bd. Can. Tech. Rep.*, 225, 1971, 18 pp.
90. Fletcher, G. L., "Accumulation of Yellow Phosphorus by Several Marine Invertebrates and Seaweed", *J. Fish. Res. Bd. Canada*, 28:793-796 (1971).
91. Peer, D. L., "Observations on Mortalities of Benthic Organisms After Contamination of the Bottom of Long Harbour, "Iacentia Bay, Newfoundland, with Elemental Phosphorus", *Circ. No. 2*, Nov, 1972, Atlantic Regional Office, Research and Development, *Fish. Res. Bd. Canada*, Halifax, N.S., pp. 181-186.

LITERATURE CITED (Cont)

92. Black, R., "Growth Rates of Intertidal Molluscs as Indicators of Effects of Unexpected Incidents of Pollution", *Fish. Res. Bd. Canada*, Biological Station, St. John's, Newfoundland, preprint courtesy of author, 1973.
93. Dyer, W. J., D. F. Hiltz, R. G. Ackman, J. Hingley, and G. L. Fletcher, "In vivo Assimilation by Cod Muscle and Liver Tissue of Elemental Phosphorus from Polluted Sea Water", *J. Fish. Res. Bd. Canada*, 27:1131-1139 (1970).
94. Ackman, R. G., R. F. Addison, and J. Hingley, "An Assessment of the Assimilation of Elemental Phosphorus by Newfoundland Marine Organisms in the 1969 Pollution Problem and in 1970 Monitoring Operations", *Fish. Res. Board Can. Tech. Rep.*, 208, 1970, 39 pp.
95. Coburn, D. R., J. B. Dewitt, J. V. Derby, Jr., and E. Ediger, "Phosphorus Poisoning in Waterfowl", *J. Am. Pharm. Assoc.*, 39: 151-158 (1950).
96. Blumbergs, P. et al., "Chemical Process Studies for Commercially Unavailable Materials", Edgewood Arsenal Contract DAAA 15-69-C-0584, Final Report, 138 pp., 1973.
97. Van Wazer, J. R., "Phosphorus and Its Compounds. Vol. I", Interscience Publishers, Inc., New York, 1958, pp. 359-366.
98. Bulen, W. A. and D. S. Frear, "Effect of Orthophosphite on Respiration, Growth, and Nitrogen Fixation of *Azotobacter vinelandii*", *Arch. Biochem. Biophys.*, 66:502-503 (1957).
99. Casida, L. E., Jr., "Microbial Oxidation and Utilization of Orthophosphite During Growth", *J. Bacteriol.*, 80:237-241 (1960).
100. Robertson, H. E. and P. D. Boyer, "The Biological Inactivity of Glucose-6-phosphite, Inorganic Phosphites and Other Phosphites", *Arch. Biochem. Biophys.*, 62:380-395 (1956).
101. Van Wazer, J. R., *op. cit.*, p. 381.
102. Van Wazer, J. R., *op. cit.*, p. 367.
103. Crewther, W. G., "The Inhibition of Formic Dehydrogenases and Formic Hydrogenlyase Systems of *Escherichia coli* by Hypophosphite", *Biochim. et Biophys. Acta*, 21:178-180 (1956).

LITERATURE CITED (Cont)

104. Grunberg-Monago, M., J. Szulmajster, and A. Prouvost, "Hydrogenlyase, Formic Dehydrogenase and Hydrogenase from *Escherichia coli*", *Compt. Rend.*, 223:1690-1692 (1951).
105. Frommel, E., A. D. Herschberg, and J. Piquet, "Effects of Inorganic Ions on the Activity of Serum Cholinesterase", *Helv. Physiol. Acta*, 2:169-191 (1944).
106. Doudoroff, P. and M. Katz, "Critical Review of Literature on the Toxicity of Industrial Wastes and Their Components to Fish. I. Alkalies, Acids and Inorganic Gases", *Sewage and Ind. Wastes*, 22:1432-1455 (1950).
107. Datsko, V. G., "Hydrochemical Conditions in the Azov Sea Under an Anoxic Regime", *Gidrokhim. Materialy*, 25:28 (1955). (C.A., 52: 6868, 1958).
108. Luning, O. and K. Brohm, "Occurrence of Phosphine in Spring Waters", *Z. Untersuch. Lebensm.*, 61:443 (1931). (C.A., 26:541, 1932).
109. Luning, O. and K. Brohm, "Occurrence of Phosphine in Spring Waters", *Z. Untersuch. Lebensm.*, 66:460 (1933). (C.A., 28:6219, 1934).
110. Hilton, H. W. and W. H. Robison, "Fate of Zinc Phosphide and Phosphine in the Soil-Water Environment", *J. Agr. Food Chem.*, 20:1209-1213 (1972).
111. Karl-kroupa, E., C. F. Callis, and E. Seifter, "Stability of Condensed Phosphates in Very Dilute Solution", *Ind. Eng. Chem.*, 49: 2061-2062 (1957).
112. Jangaard, P. M., "The Role Played by the Fisheries Research Board of Canada in the 'Red' Herring Phosphorus Pollution Crisis in Placentia Bay, Newfoundland", Circ. No. 1, Office of the Atlantic Regional Director, Research, Fish. Res. Bd. Canada, Halifax, N. S., 1970, 20 pp.
113. Odense, P. H., Y. M. Marbougali, T. C. Leung, and V. H. Logan, "Hematological and Histological Studies Resulting from the Pollution Problem at Long Harbour, Placentia Bay, Newfoundland", Circ. No. 2, Nov. 1971, Atlantic Regional Office, Research and Development, Fish. Res. Bd. Canada, Halifax, N. S., pp. 195-222.

LITERATURE CITED (Cont)

114. Mazmanidi, H. D., "Pathomorphology of the Phosphorus Poisoning of Fish", *Vop. Vod. Toksikol.*, 128 (1970). (C.A., 74:73450, 1971).
115. Fletcher, G. L., "The Acute Toxicity of a Yellow Phosphorus Contaminated Diet to Brook Trout", *Bull. Environ. Contam. Toxicol.*, 10(2):123-128 (1973).
116. Aiken, D. E. and E. H. Byard, "Histological Changes in Lobsters (*Homarus americanus*) Exposed to Yellow Phosphorus", *Science*, 176:1434-1435 (1972).
117. Dyer, W. J., D. F. Hiltz, R. G. Ackman, J. Hingley, G. L. Fletcher, and R. F. Addison, "Stability of Elemental Phosphorus in Edible Muscle Tissue of Cod During Processing, Including Icing, Freezing and Thawing, Frozen Storage, Salting and Cooking", *J. Fish. Res. Bd. Canada*, 29:1053-1060 (1972).
118. Anon., General Technologies Corp., "Development Document for Effluent Limitations Guidelines and Standards of Performance: Non-Fertilizer Phosphorus Chemicals Industry", US Environmental Protection Agency Contract No. 68-01-1513, June 1973; letter from K. A. Booman, Technical Director, The Soap and Detergent Association, to the Chemical and Environmental Research Committee members.
119. Addison, R. F., "Analysis of Elemental Phosphorus and Some of Its Compounds by Gas Chromatography", *Proc. Int. Symp. Identification Measurement Environmental Pollutants*, Ottawa, Ont., June 14-17, 1971, pp. 386-390.
120. Bullock, E. and M. J. Newlands, "Decomposition of Phosphorus in Water", *Proc. Conf. Poll. Chem. Inst. Can.*, Halifax, N. S., Aug 24-26, 1969, pp. 23-24.
121. Anon., "Oldbury Electro-Chemical's Phosphorus Plant Solves Problem of Waste Disposal by Installing Unit for Treatment of Effluent with Chlorine", *Chem. Eng. News*, 30:466 (1952).
122. Barber, J. C., "Waste Effluent; Treatment and Reuse", *Chem. Eng. Prog.*, 65:70-73 (1969).
123. Ackman, R. G. and R. F. Addison, "Direct Determination of Elemental Phosphorus by Gas-Liquid Chromatography", *Proc. Conf. Pollut.*, St. Mary's Univ., Halifax, N. S., Aug 24-26, 1969, pp. 140-147.

LITERATURE CITED (Cont)

124. Addison, R. F. and R. G. Ackman, "Direct Determination of Elemental Phosphorus by Gas-Liquid Chromatography", *J. Chromatog.*, 47:421-426 (1970).
125. Addison, R. F., R. G. Ackman, L. J. Cowley, D. Mascaluk, and S. Pond, "Elemental Phosphorus Levels in Water Samples from Long Harbour, Newfoundland (July 1969-June 1970)", *Fish. Res. Board Can. Tech. Rep.*, 254, 1971, 29 pp.
126. Addison, R. F., M. E. Zinck, R. G. Ackman, P. Chamut, and A. Jamieson, "Results of the Elemental Phosphorus Monitoring Program, Long Harbour, Newfoundland (July 1970-April 1971)", *Fish. Res. Board Can. Tech. Rep.*, 203, 1972, 25 pp.
127. Ackman, R. G., R. F. Addison, P. J. Ke, and J. C. Sipos, "Examination of Bottom Deposits, Long Harbour, Newfoundland, for Elemental Phosphorus and for Fluorides", *Fish. Res. Board Can. Tech. Rep.*, 233, 1971, 18 pp.
128. Bohl, C. D. and E. F. Kaelble, "Determination of Elemental Phosphorus in Air by Gas chromatography", *Am. Ind. Hyg. Assoc. Conf.*, May 14-19, 1972, San Francisco, CA.
129. Gorzny, K., "Analysis of Elemental Phosphorus" in "Analytical Chemistry of Phosphorus Compounds", M. Halmann, Editor, *Interscience*, New York, 1972, pp. 361ff.
130. Fetchin, J. A. and M. Grayson, "Tervalent Phosphorus Compounds" in "Analytical Chemistry of Phosphorus Compounds", M. Halmann, Editor, *Interscience*, New York, 1972, pp. 371-383.
131. Dechant, R., G. Sanders, and R. Gaul, "Determination of Phosphorus in Air", *Am. Ind. Hyg. Assoc. J.*, 27:75-78 (1966).
132. Berck, B., W. E. Westlake, and F. A. Gunther, "Microdetermination of Phosphine by Gas-Liquid chromatography with Microcoulometric, Thermionic, and Flame Photometric Detection", *J. Agr. Food Chem.*, 18:143-147 (1970).
133. Van Wazer, J. R., *op. cit.*, pp. 415-418.

LITERATURE CITED (Cont)

134. Ohashi, S., "Phosphorus Oxides and Oxo Acids" in "Analytical Chemistry of Phosphorus Compounds", M. Halmann, Editor, *Interscience*, New York, 1972, pp. 409ff.
135. Seiler, H., "Thin Layer Chromatography of Inorganic Ions" in "Thin Layer Chromatography", E. Stahl, Editor, *Academic Press*, New York, 1965, pp. 469-483.
136. Butts, W. C., "Gas Chromatography of Trimethylsilyl Derivatives of Common Anions", *Anal. Letters*, 3:29-34 (1970).

APPENDIX A
SOURCES CONSULTED

A. Books:

1. Christensen, H. E. (ed.): "Toxic Substances List". 1972 Edition. USDHEW/NIOSH, 1972.
2. Fishbein, L., W. G. Flamm, and H. L. Falk: "Chemical Mutagens". New York and London, Academic Press, 1970.
3. Gafafer, W. M. (ed.): "Occupational Disease". USDHEW, 1964.
4. Gleason, M. N., R. E. Gosselin, H. C. Hodge, and R. P. Smith: "Clinical Toxicology of Commercial Products. Acute Poisoning". Third Edition. The Williams and Wilkins Co., Baltimore, 1969.
5. Goodman, L. S. and A. Gilman: "The Pharmacological Basis of Therapeutics". Fourth Edition. The Macmillan Company, New York, 1970.
6. Hamilton, A. and H. L. Hardy: "Industrial Toxicology". Second Edition. P. B. Hoeber, Inc., New York, 1949.
7. Hartwell, J. L.: "Survey of Compounds Which have been Tested for Carcinogenic Activity". Second Edition. 1951.
Supplement 1 (Shubik and Hartwell), 1957.
Supplement 2 (Shubik and Hartwell), 1969.
8. Hayes, W. J., Jr.: "Clinical Handbooks on Economic Poisons". USDHEW/PHS, Communicable Disease Center, Atlanta, GA, 1963.
9. International Labour Office: "Occupation and Health (Encyclopedia of Hygiene, Pathology and Social Welfare)". Vol II (I-Z), Geneva, 1934.
10. Negherbon, W. O.: "Handbook of Toxicology. Vol III. Insecticides A Compendium". W. B. Saunders Co., Philadelphia, 1959.
11. Patty, F. A., D. W. Fassett, and D. D. Irish (eds.): "Industrial Hygiene and Toxicology, Vol. 2. Toxicology". Interscience Publishers, New York, 1963.
12. Sax, N. I.: "Dangerous Properties of Industrial Materials". Third Edition. Reinhold Publishing Corp., New York, 1968.

APPENDIX A (Cont)

13. Spector, W. S. (ed.): "Handbook of Toxicology, Vol. 1. Acute Toxicities of Solids, Liquids and Gases to Laboratory Animals". W. B. Saunders Co., Philadelphia, 1956.
14. Stecher, P. G (ed.): "The Merck Index (of Chemicals and Drugs)". Eighth Edition. Merck and Co., Inc., Rahway, New Jersey, 1968.
15. Sutton, H. E. and M. I. Harris (eds.): "Mutagenic Effects of Environmental Contaminants". Academic Press, New York and London, 1972.
16. "Water Quality Criteria Data Book". Selected Data from the Literature thru 1968, Water Pollution Control Research Series, EPA, Volumes 1-4, 1970-1972.
17. Williams, R. T.: "Detoxication Mechanisms (The Metabolism and Detoxication of Drugs, Toxic Substances and Other Organic Compounds)". Second Edition. John Wiley and Sons, Inc., New York, 1959.

B. Journals Searched for Phosphorus:

1. *Chemical Abstracts*, from Vol. 1, 1907 to data.

Includes 1st decennial index, 1-10 (1907-1916)

2nd " " 11-20 (1917-1926)

3rd " " 21-30 (1927-1936)

4th " " 31-40 (1937-1946)

5th " " 41-50 (1947-1956)

6th collective " 51-55 (1957-1961)

7th " " 56-65 (1962-1966)

8th " " 66-75 (1967-1971)

Vols. 76, 77 (1972), and 78 (1973) and 79 (1973) to date.

[Key words: phosphorus, white phosphorus]

2. *Biological Abstracts*, from Vol. 1, 1927 to date (1973, 55).

[Key words: phosphorus, white]

3. *Toxicity Bibliography*, from Vol. 1, 1968 to date (Vol. 6, 1973).

[Key words: phosphorus, white]

APPENDIX A (Cont)

C. Computer Searches of Literature for Phosphorus:

1. Defence Documentation Center, Defense Supply Agency. Cameron Agency, Alexandria, VA 22314, 1940 to date.

[Key words: phosphorus, white phosphorus, phossy water, plasticized white phosphorus].

(See Search Control No. 002787)

2. National Library of Medicine: Medline retrieval file (1969 to date) and Toxicon (1966 to date).

[Key words: phosphorus, white]

APPENDIX B

UNCITED REFERENCES

A. Relevant References not Quoted in Text

1. Chamut, P. S., S. G. Pond, and V. R. Taylor, "The 'Red Herring' Pollution Crisis in Placentia Bay, Newfoundland. A General Description and Chronology", Circ. No. 2, Nov 1972, Atlantic Regional Office, Research and Development, Fish. Res. Bd. Canada, Halifax, N. S., pp. 29-43.

2. Cornick, J. W., R. M. Mackelvie, and J. E. Stewart, "A Note on Bacteriological Examination of Moribund Herring from Placentia Bay", *ibid.*, p. 53.

3. Griffith, E. J., A. Beeton, J. M. Spencer, and D. T. Mitchell, (Ed.), "Environmental Phosphorus Handbook", Wiley-Interscience, New York, 1973, 718 pp.

4. Hodder, V. M., L. S. Parsons, and J. H. C. Pippy, "The Occurrence and Distribution of 'Red' Herring in Placentia Bay, February - April 1969", Circ. No. 2, Nov 1972, Atlantic Regional Office, Research and Development Fish. Res. Bd. Canada, Halifax, N. S., pp. 45-52.

5. Li, M. F., G. S. Traxler, and L. M. Langille, "Toxic Effect of Elemental Phosphorus on L-cells Cultivated in Suspension", *Can. J. Zool.*, 48:133 (1969).

6. Pippy, J. H. C., V. M. Hodder, and L. S. Parsons, "Symptoms of 'Red' Herring in Relation to Mass Mortalities in Placentia Bay, February-April 1969", Circ. No. 2, Nov 1972, Atlantic Regional Office, Research and Development, Fish. Res. Bd. Canada, Halifax, N. S., pp. 187-191.

7. Trites, R. W., "Capacity of an Estuary to Accept Pollutants", *Proc. Conf. Poll. Chem. Inst. Can.*, Halifax, N. S., Aug. 24-26, 1969, pp. 23-24.

B. References Not Considered Relevant

1. Griffith, E. J. and M. Grayson, "Topics in Phosphorus Chemistry", Vol. 1-7, Interscience, New York, 1964-1972. (Does not cover elemental phosphorus).

APPENDIX B (Cont)

2. MacKay, E. M. and J. Oliver, "Renal Damage Following the Ingestion of a Diet Containing an Excess of Inorganic Phosphate", *J. Exp. Med.*, 61: 319-333 (1935).

3. MacKay, E. M. and J. Oliver, "Renal Damage Following the Administration of Inorganic Phosphate", *Proc. Soc. Exp. Biol. Med.*, 28:234-235 (1930).

4. Dalhamn, T. and M. L. BoHolma, "Animal Experiments on the Spread of Inhaled Radioactive Red Phosphorus in the Body", *Arch. Ind. Health*, 20:429-431 (1959).

C. References Requiring Translation

1. Oddo and Olmer, "Recherches sur l'intoxication phosphorée expérimentale", *Comp. Rend. Soc. Biol.*, 56:901-902 (1904).

2. Neubauer, E. and O. Porges, "Über Nebenniereninsuffizienz bei Phosphorvergiftung", *Biochem. Z.*, 32:290-307 (1911).

3. Leo, H., "Über das Wesen der Organverfettungen nach Phosphorvergiftung", *Biochem. Z.*, 48:297-301 (1913).

4. Ivančević, I., "Über experimentelle Phosphorvergiftung mit besonderer Berücksichtigung des Blutzuckers", *Arch. Exp. Pathol. Pharmacol.*, 122:24-43 (1927).

5. Zernik, F., "Beiträge zur Kenntnis der perkutanen Phosphorwirkung", *Z. Exp. Med.*, 70:1-15 (1930).

6. Hennigues, V. and U. Klausen, "Untersuchungen über den Serumalbumin- und Serumglobulin-gehalt des Serums unter wechselnden Umständen", *Biochem. Z.*, 254:414-433 (1933).

7. Hurka, W., "Tierexperimentelle Untersuchungen mit weissem Phosphor und seinen Oxydationsprodukten", *Klin. Med. Wien.*, 2:1014-1024 (1947).

8. Venturi, V. M., "Le frazioni dello zolfo urinario nell'intossicazione sperimentale da fosforo", *Boll. Soc. Ital. Biol. Sper.*, 24:92-94 (1948).

9. Venturi, V. M., "Azione della metionina sul ricambio solforato e sulla solfoconjugazione in animali normali o sperimentalmente intossicati con fosforo", *Arch. Int. Pharmacodyn.*, 83:165-173 (1950).

APPENDIX B (Cont)

10. Lattes, L., "Über den Fettgehalt des Blutes des Hundes unter normalen und unter verschiedenen experimentellen Verhältnissen (Verdauung, Hungern, Phosphor-, Phloridzin- und Chloroformvergiftung)", *Arch. Exp. Pathol. Pharmacol.*, 66:132-142 (1911).
11. Wordell, H., "Vergleichende Untersuchungen über den zeitlichen Verlauf der Phosphorausscheidung im Urin Gesunder und Nierenkranken", *Z. Exp. Med.*, 42:389-399 (1924).
12. Gubser, A., "Zur Chemie der Leberverfettung", *Biochem. Z.*, 198:65-80 (1928).
13. Loewy, A. and G. Cronheim, "Weitere Beiträge zur Chemie der Leberverfettung", *Biochem. Z.*, 257:267-288 (1933).
14. Katano, S., "Studien ueber Blut und Harn beim experimentellen Stauungsikterus. II. Mitteilung: Untersuchungen bei der Phosphor oder Arsen-vergiftungen", *J. Kumamoto Med. Soc.*, 10:177-178 (1934).
15. Nakamura, T., "Experimentelle Untersuchungen über die akute Phosphorvergiftung. II. Mitteilung. Zeitliche Beobachtung der Nierenveränderung", *Okayama-Igakkai-Zasshi*, 46:2804-2805 (1934).
16. Kawada, T. and F. Yamasaki, "Zur Kenntnis des Phosphorharns beim Menschen", *J. Biochem. (Japan)* 21:423-425 (1935).
17. Fischler, F. and K. E. Schulte, "Ueber die Entgiftung tödlicher Phosphorgaben durch Stoffwechselbeeinflussung der Leber", *Munch. Med. Wochschr.*, 88:854-857 (1941).
18. Guccione, F., "Il SRE nelle tossicosi da fosforo, pulegone, arsenico negli animali interi e smilzati", *Hematologica*, 10:63-86 (1929).
19. Filippi, E. and S. Oberto, "Osservazioni sopra alcuni metodi recenti per la ricerca tossicologica del Fosforo", *Arch. Farmacol. Sper.*, 8:211-219 (1909).
20. Rettig, H., "Zur Frage des toxogenen Eiweisszerfalls bei der Phosphorvergiftung", *Arch. Exp. Pathol. Pharmacol.*, 76:345-366 (1914).
21. Heinlein, H. and M. Angermann, "Morphologische und Chemische Blutveränderungen bei Phosphorvergiftung. Untersuchungen am Phosphorhund", *Z. Allg. Pathol.*, 58:81-95 (Supplement) (1933).

APPENDIX B (Cont)

22. Láng, S. and I. Franreisz, "Über den Einfluss der Chronischen Phosphor- und Arsenvergiftung auf die Plasmaeiweisskörper", *Z. Ges. Exp. Med.*, 89:124-130 (1933).

23. Mariconda, P., "Sull'Intossicazione cronica da Fosforo-recerche sperimentali", *Arch. Farmacol. Sper.*, 39:24-33 (1925).

24. Starkenstein, E., "Zur experimentellen analyse Chronischer Vergiftungen", *Klin. Wochenschr.*, 11:1697-1702 (1932).

25. Michaelis, P., "Das Wesen der Chronischen Phosphorwirkung und der Phosphornekrose", *Arch. Gewerbepathol. Hyg.*, 7:477-485 (1936).